



Perfect 3-colorings of some generalized peterson graph

Mehdi Alaeiyan and Zahra Shokoohi¹

School of Mathematics, Iran University of Science and Technology, Narmak, Tehran 16846, Iran

Abstract

The notion of a perfect coloring, introduced by Delsarte, generalizes the concept of completely regular code. A perfect z -colorings of a graph is a partition of its vertex set. It splits vertices into z parts P_1, \dots, P_z such that for all $i, j \in \{1, \dots, z\}$, each vertex of P_i is adjacent to p_{ij} vertices of P_j . The matrix $P = (p_{ij})_{i,j \in \{1, \dots, z\}}$, is called parameter matrix. In this article, we classify all the realizable parameter matrices of perfect 3-colorings of some the generalized peterson graph.

Keywords: Parameter matrices, Perfect coloring, Equitable partition, Generalized peterson graph.

Mathematical Subject Classification 03E02, 05C15, 68R05

1 Introduction

The concept of a perfect z -coloring plays a significant role in graph theory, algebraic combinatorics, and coding theory (completely regular codes). There is another phrase for this concept in the writing as “equitable partition” (see [9]). In 1973, Delsarte conjectured the non-existence of nontrivial perfect codes in Johnson graphs. Since then, some effort has been made to count the parameter matrices of some Johnson graphs, including $J(4, 2)$, $J(5, 2)$, $J(6, 2)$, $J(6, 3)$, $J(7, 3)$, $J(8, 3)$, $J(8, 4)$, and $J(v, 3)$ (v odd) (see [3, 4, 8]).

Fon-Der-Flass count the parameter matrices (perfect 2-colorings) of n -dimensional hypercube Q_n for $n < 24$. He also obtained some constructions and a necessary condition for the existence of perfect 2-colorings of the n -dimensional cube with a given parameter matrix (see[5, 6, 7]). In this article, we classify the parameter matrices of all perefect 3-colorings of some generalized peterson graph.

Some generalized peterson graph including $GP(7, 1)$, $GP(8, 1)$, $GP(8, 2)$ and $GP(8, 3)$ given as follow:

¹speaker




Figure 1: Some generalized peterson graph

Definition 1.1. The generalized peterson graph $GP(n, k)$ has vertices, respectively, edges given by

$$V(GP(n, k)) = \{a_i, b_i : 0 \leq i \leq n - 1\},$$

$$E(GP(n, k)) = \{a_i a_{i+1}, a_i b_i, b_i b_{i+k} : 0 \leq i \leq n - 1\},$$

Where the subscripts are expressed as integers modulo $n (\geq 5)$, and $k (\geq 1)$ is the skip.

Definition 1.2. For a graph G and an integer z , a mapping $T : V(G) \rightarrow \{1, 2, \dots, z\}$ is called a perfect z -coloring with matrix $P = (p_{ij})_{i,j \in \{1, \dots, z\}}$, if it is surjective, and for all i, j , for every vertex of color i , the number of its neighbours of color j is equal to p_{ij} . The matrix P is called the parameter matrix of a perfect coloring. In the case $z = 3$, we call the first color white that show by W , the second color black that show by B and the third color red that show by R . In this article, we generally show a parameter matrix by

$$P = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}.$$

Remark 1.3. In this paper, we consider all perfect 3-colorings, up to renaming the colors; i.e. We identify the perfect 3-coloring with the matrices

$$\begin{bmatrix} d & c & b \\ g & i & h \\ d & e & f \end{bmatrix}, \begin{bmatrix} e & d & f \\ b & a & c \\ h & g & i \end{bmatrix}, \begin{bmatrix} e & f & d \\ h & i & g \\ b & c & a \end{bmatrix}, \begin{bmatrix} i & h & g \\ f & e & d \\ c & b & a \end{bmatrix}, \begin{bmatrix} i & g & h \\ c & a & b \\ f & d & e \end{bmatrix}.$$

Obtained by switching the colors with original coloring .

2 Preliminaries

In this section, we present some results concerning necessary conditions for the existence of perfect 3-colorings of the generalized peterson graph of $GP(7,1)$, $GP(8,1)$, $GP(8,2)$ and $GP(8,3)$ with a given parameter matrix

$$P = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

The simplest necessary condition for the existence of perfect 3-colorings of the generalized peterson

$$a + b + c = d + e + f = g + h + i = 3.$$

By using this condition and some computation, it is clear that we should consider 18 matrices .These matrices are listed below:

$$\begin{aligned} P_1 &= \begin{bmatrix} 0 & 0 & 3 \\ 0 & 0 & 3 \\ 1 & 1 & 1 \end{bmatrix}, & P_2 &= \begin{bmatrix} 0 & 0 & 3 \\ 0 & 0 & 3 \\ 1 & 2 & 0 \end{bmatrix}, & P_3 &= \begin{bmatrix} 0 & 0 & 3 \\ 0 & 1 & 2 \\ 1 & 1 & 1 \end{bmatrix}, & P_4 &= \begin{bmatrix} 0 & 0 & 3 \\ 0 & 2 & 1 \\ 1 & 2 & 0 \end{bmatrix}, \\ P_5 &= \begin{bmatrix} 0 & 0 & 3 \\ 0 & 2 & 1 \\ 1 & 1 & 1 \end{bmatrix}, & P_6 &= \begin{bmatrix} 0 & 0 & 3 \\ 0 & 2 & 1 \\ 2 & 1 & 0 \end{bmatrix}, & P_7 &= \begin{bmatrix} 0 & 1 & 2 \\ 1 & 0 & 2 \\ 1 & 1 & 1 \end{bmatrix}, & P_8 &= \begin{bmatrix} 0 & 1 & 2 \\ 1 & 1 & 1 \\ 2 & 1 & 0 \end{bmatrix}, \\ P_9 &= \begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 0 \\ 1 & 0 & 2 \end{bmatrix}, & P_{10} &= \begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 0 \\ 2 & 0 & 1 \end{bmatrix}, & P_{11} &= \begin{bmatrix} 0 & 3 & 0 \\ 1 & 0 & 2 \\ 0 & 1 & 2 \end{bmatrix}, & P_{12} &= \begin{bmatrix} 1 & 0 & 2 \\ 0 & 0 & 3 \\ 1 & 2 & 0 \end{bmatrix}, \\ P_{13} &= \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \\ 1 & 1 & 1 \end{bmatrix}, & P_{14} &= \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \\ 1 & 2 & 0 \end{bmatrix}, & P_{15} &= \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 1 & 1 & 1 \end{bmatrix}, & P_{16} &= \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}, \\ P_{17} &= \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 0 \\ 1 & 0 & 2 \end{bmatrix}, & P_{18} &= \begin{bmatrix} 1 & 2 & 0 \\ 1 & 0 & 2 \\ 0 & 1 & 2 \end{bmatrix}. \end{aligned}$$

Theorem 2.1. [9] *If T is a perfect coloring of a graph G in z colors, then any eigenvalue of T is an eigenvalue of G .*

Theorem 2.2. [1] *Suppose that T is a perfect 3- coloring with matrix $\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$, in the connected graph G .Then in this case, none of the following situations will occur.*

- (1) $b = c = 0$,
- (2) $d = f = 0$,
- (3) $g = h = 0$,
- (4) $b = 0 \leftrightarrow d = 0, c = 0 \leftrightarrow g = 0, h = 0 \leftrightarrow f = 0$.

Theorem 2.3. [2] *Let T a perfect 3-coloring of a graph G with matrix $P = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$.*

(1) If $b, c, f \neq 0$, then

$$|W| = \frac{|V(G)|}{\frac{b}{d} + 1 + \frac{c}{g}}, \quad |B| = \frac{|V(G)|}{\frac{d}{b} + 1 + \frac{f}{h}}, \quad |R| = \frac{|V(G)|}{\frac{h}{f} + 1 + \frac{g}{c}}.$$

(2) If $b = 0$, then

$$|W| = \frac{|V(G)|}{\frac{c}{g} + 1 + \frac{ch}{fg}}, \quad |B| = \frac{|V(G)|}{\frac{f}{h} + 1 + \frac{fg}{ch}}, \quad |R| = \frac{|V(G)|}{\frac{h}{f} + 1 + \frac{g}{c}}.$$

(3) If $c = 0$, then

$$|W| = \frac{|V(G)|}{\frac{b}{d} + 1 + \frac{bf}{dh}}, \quad |B| = \frac{|V(G)|}{\frac{d}{b} + 1 + \frac{f}{h}}, \quad |R| = \frac{|V(G)|}{\frac{h}{f} + 1 + \frac{dh}{bf}}.$$

(4) If $f = 0$, then

$$|W| = \frac{|V(G)|}{\frac{b}{d} + 1 + \frac{c}{g}}, \quad |B| = \frac{|V(G)|}{\frac{d}{b} + 1 + \frac{cd}{bg}}, \quad |R| = \frac{|V(G)|}{\frac{g}{c} + 1 + \frac{bg}{cd}}.$$

Theorem 2.4. [1] If $P = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$ be a parameter matrix of a k -regular graph, then the eigenvalues of P are

$$\lambda_{1,2} = \frac{\text{tr}(P) - k}{2} \pm \sqrt{\left(\frac{\text{tr}(P) - k}{2}\right)^2 - \frac{\det(P)}{k}}, \quad \lambda_3 = k.$$

Remark 2.5. The distinct eigenvalues of the graph $GP(7, 1)$ are the numbers 3, 1, The distinct eigenvalues of graph $GP(8, 1)$ are the numbers 3, 1, -1 , The distinct eigenvalues of graph $GP(8, 2)$ are the numbers 1, 3 and the distinct eigenvalues of graph $GP(8, 3)$ are the numbers 3, 1, -1 .

3 Perfect 3- colorings of some generalized peterson graph

The parameter matrices of $GP(7, 1)$, $GP(8, 1)$, $GP(8, 2)$ and $GP(8, 3)$ graphs are enumerated in the next theorems.

Theorem 3.1. The graph $GP(7, 1)$ has no perfect 3-colorings.

Proof. A parameter matrix corresponding to perfect 3-colorings of the graph $GP(7, 1)$ may be one of the matrices P_1, \dots, P_{18} . By using Theorem 2.1 and Theorem 2.4, we can see that only the matrices $P_3, P_4, P_5, P_6, P_{10}, P_{12}, P_{15}$, and P_{18} can be a parameter matrices. By using Theorem 2.3, matrices P_3, P_6, P_{10}, P_{12} , and P_{15} cannot be a parameter matrices, because the number of white, black and red, are not an integer. For matrix P_4 , each vertex with color white has three adjacent vertices with color red. Now have the following possibilities:

- (1) $T(a_1) = T(a_{11}) = W$, $T(a_3) = T(a_5) = T(a_9) = T(a_{10}) = B$, $T(a_2) = T(a_4) = T(a_7) = T(a_8) = T(a_{12}) = R$ then $T(a_2) = T(a_{13}) = T(a_{14}) = B$, which is a contradiction with the second row of matrix P_4 .
- (2) $T(a_1) = T(a_8) = T(a_9) = T(a_{14}) = B$, $T(a_3) = W$, $T(a_2) = T(a_4) = T(a_6) = T(a_7) = T(a_{10}) = T(a_{13}) = R$ then $T(a_5) = T(a_{11}) = T(a_{12}) = B$, which is a contradiction with the second row of matrix P_4 . Hence graph $GP(7, 1)$ has no perfect 3-colorings with matrix P_4 .

Similar to matrix P_4 , we proof for matrix P_5 and P_{18} as follows:

For matrix P_5 , each vertex with color white has three adjacent vertices with color red. Now have the following possibilities:

- (3) $T(a_3) = T(a_6) = W$, $T(a_7) = T(a_8) = T(a_{12}) = T(a_{14}) = B$, $T(a_2) = T(a_4) = T(a_5) = T(a_9) = T(a_{13}) = R$ then $T(a_4) = T(a_{10}) = R$, $T(a_{11}) = B$ which is a contradiction with the second row of matrix P_5 .
- (4) $T(a_1) = T(a_2) = T(a_5) = T(a_6) = T(a_{12}) = B$, $T(a_{10}) = W$, $T(a_3) = T(a_4) = T(a_8) = T(a_9) = T(a_{11}) = T(a_{13}) = R$ then $T(a_7) = B$ and $T(a_{14}) = W$, which is a contradiction with the second row of matrix P_5 . Hence graph $GP(7, 1)$ has no perfect 3-colorings with matrix P_5 .

For matrix P_{18} , each vertex with color white has two adjacent vertices with color black. Now have the following two possibilities:

- (5) $T(a_1) = T(a_2) = T(a_3) = T(a_4) = T(a_5) = T(a_8) = T(a_{12}) = T(a_{14}) = R$, $T(a_7) = T(a_9) = T(a_{11}) = B$, $T(a_6) = W$ then $T(a_{13}) = R$, which is a contradiction with the second three of matrix P_{18} .
- (6) $T(a_1) = T(a_4) = T(a_5) = T(a_9) = T(a_{10}) = T(a_{12}) = T(a_{14}) = R$, $T(a_2) = T(a_3) = B$, $T(a_8) = T(a_{11}) = W$ then $T(a_6) = T(a_7) = T(a_{13}) = B$, which is a contradiction with the second row of matrix P_{18} . Hence graph $GP(7, 1)$ has no perfect 3-colorings with matrix P_{18} .

□

Theorem 3.2. *The graph $GP(8, 1)$ has a perfect 3-colorings with the matrices P_7 and P_{13} .*

Proof. A parameter matrix corresponding to perfect 3-colorings of the graph $GP(8, 1)$ may be one of the matrices P_1, \dots, P_{18} . Using the Theorems 2.1 and 2.4 matrices $P_3, P_4, P_5, P_6, P_7, P_{10}, P_{12}, P_{13}, P_{15}, P_{16}$ and P_{18} can be a parameter matrices. By using Theorem 2.3 matrices P_5, P_6, P_7 and P_{13} cannot be a parameter matrices, because of the number of white colors is not integer.

Consider the mapping T_1 and T_2 as follows:

$$\begin{aligned}
 T_1(a_1) &= T_1(a_5) = T_1(a_{11}) = T_1(a_{15}) = W, & T_1(a_2) &= T_1(a_6) = T_1(a_{12}) = T_1(a_{16}) = B, \\
 T_1(a_3) &= T_1(a_4) = T_1(a_7) = T_1(a_8) = T_1(a_9) = T_1(a_{10}) = T_1(a_{13}) = T_1(a_{14}) = R. \\
 \\
 T_2(a_2) &= T_2(a_3) = T_2(a_6) = T_2(a_7) = W, & T_2(a_9) &= T_2(a_{12}) = T_2(a_{13}) = T_2(a_{16}) = B, \\
 T_2(a_1) &= T_2(a_4) = T_2(a_5) = T_2(a_8) = T_2(a_{10}) = T_2(a_{11}) = T_2(a_{14}) = T_2(a_{15}) = R.
 \end{aligned}$$

It is clear that T_1 and T_2 are perfect 3-coloring with the matrices P_7 and P_{13} respectively. □

Theorem 3.3. *The graph $GP(8, 2)$ has no perfect 3-colorings.*

Proof. A parameter matrix corresponding to perfect 3-colorings of the graph $GP(8, 2)$ may be one of the matrices P_1, \dots, P_{18} . By using Theorem 2.1 and Theorem 2.4, we can see that only the matrices $P_3, P_4, P_5, P_6, P_{10}, P_{12}, P_{15}$ and P_{18} can be a parameter matrices. By using Theorem 2.3, matrices $P_4, P_5, P_{10}, P_{12}, P_{15}, P_{18}$ cannot be a parameter matrices, because the number of white, black and red, are not an integer. For matrix P_6 , each vertex with color white has three adjacent vertices with color red. Now have the following possibilities:

- (1) $T(a_1) = T(a_5) = T(a_7) = T(a_8) = T(a_{10}) = T(a_{11}) = T(a_{14}) = T(a_{15}) = R$, $T(a_2) = T(a_3) = B$, $T(a_6) = T(a_9) = T(a_{13}) = T(a_{16}) = W$ then $T(a_4) = B$ and $T(a_{12}) = W$, which is a contradiction with the second row of matrix P_6 .
- (2) $T(a_1) = T(a_5) = T(a_{11}) = T(a_{15}) = R$, $T(a_2) = T(a_9) = T(a_{12}) = T(a_{13}) = W$, $T(a_7) = T(a_8) = T(a_{14}) = T(a_{16}) = B$ then $T(a_3) = T(a_4) = R$, which is a contradiction with the three row of matrix P_6 . Hence graph $GP(8, 2)$ has no perfect 3-colorings with the matrix P_6 .

□

Theorem 3.4. *The graph $GP(8, 3)$ has no perfect 3-colorings.*

Proof. A parameter matrix corresponding to perfect 3-colorings of the graph $GP(8, 3)$ may be one of the matrices P_1, \dots, P_{18} . By using Theorem 2.1 and Theorem 2.4, we can see that only the matrices $P_3, P_4, P_5, P_6, P_7, P_{10}, P_{12}, P_{13}, P_{15}$ and P_{18} can be a parameter matrices. By using Theorem 2.3, matrices $P_3, P_4, P_5, P_{10}, P_{12}, P_{14}, P_{15}$, and P_{18} cannot be a parameter matrices, because the number of white, black and red, are not an integer. For matrix P_6 , each vertex with color white has three adjacent vertices with color red. Now have the following possibilities:

- (1) $T(a_1) = T(a_3) = T(a_9) = T(a_{13}) = T(a_{14}) = R, T(a_4) = T(a_{10}) = T(a_{11}) = T(a_{15}) = T(a_{16}) = B, T(a_2) = T(a_8) = T(a_{12}) = W$ then $T(a_5) = B$ and $T(a_6) = T(a_7) = R$, which is a contradiction with the three row of matrix P_6 .
- (2) $T(a_1) = T(a_{11}) = T(a_{13}) = W, T(a_3) = T(a_4) = T(a_7) = T(a_8) = T(a_{15}) = B, T(a_2) = T(a_5) = T(a_6) = T(a_{10}) = T(a_{12}) = T(a_{16}) = R$ then $T(a_9) = T(a_{14}) = R$, which is a contradiction with the three row of matrix P_6 . Hence graph $GP(8, 3)$ has no perfect 3-colorings with the matrix P_6 .

Similar to matrix P_6 , we can proof for the matrix P_7 as follows:

For matrix P_7 each vertex with color white has two adjacent vertices with color red. Now have the following two possibilities:

- (3) $T(a_1) = T(a_4) = T(a_5) = T(a_8) = T(a_9) = T(a_{16}) = R, T(a_3) = T(a_6) = T(a_{10}) = T(a_{15}) = B, T(a_2) = T(a_{14}) = W$ then $T(a_7) = T(a_{11}) = W$ and $T(a_{12}) = T(a_{13}) = R$, which is a contradiction with the three row of matrix P_7 .
- (4) $T(a_1) = T(a_4) = T(a_9) = T(a_{12}) = W, T(a_5) = T(a_8) = T(a_{13}) = B, T(a_2) = T(a_3) = T(a_6) = T(a_7) = T(a_{10}) = T(a_{11}) = T(a_{14}) = R$ then $T(a_{15}) = T(a_{16}) = B$, which is a contradiction with the two row of matrix P_7 . Hence graph $GP(8, 3)$ has no perfect 3-colorings with matrix P_7 .

□

Finally, we summarize the results of this paper in the following table.

Table 1: Parameter matrices of some generalized peterson graph

Graphs	Parameter Matrices
graph $GP(7,1)$	×
graph $GP(8,1)$	P_7, P_{13}
graph $GP(8,2)$	×
graph $GP(8,3)$	×

References

- [1] M. Alaeiyan and A. Mehrabani, Perfect 3-colorings of cubic graphs of order 10 , *Electronic Journal of Graph Theory and Applications (EJGTA)*,5(2) (2017),PP.194-206.
- [2] M. Alaiyan, A. Mehrabani, Perfect 3- colorings of the platonic graph, *Iranian J. Sci. Technol., Trans. A Sci.* (2017) 1-9.
- [3] S.V. Avgustinovich and I. Yu. Mogilnykh, Perfect 2-colorings of Johnson graphs $J(6, 3)$ and $J(7, 3)$, *Lecture Notes in Computer Science* 5228 (2008), 11-19.
- [4] S.V. Avgustinovich and I. Yu. Mogilnykh, Perfect colorings of the Johnson graphs $J(8, 3)$ and $J(8, 4)$ with two colors, *Journal of Applied and Industrial Mathematics* 5 (2011), 19-30.
- [5] D.G. Fon-Der-Flaass, A bound on correlation immunity, *Siberian Electronic Mathematical Reports Journal* 4 (2007), 133-135.

-
- [6] D.G. Fon-Der-Flaass, Perfect 2-colorings of a hypercube, *Siberian Mathematical Journal* 4(2007), 923-930.
- [7] D.G. Fon-der-Flaass, Perfect 2-colorings of a 12-dimensional Cube that achieve a bound of correlation immunity, *Siberian Mathematical Journal* 4 (2007), 292-295.
- [8] A.L. Gavriyuk and S.V. Goryainov, On perfect 2-colorings of Johnson graphs $J(v, 3)$, *Journal of Combinatorial Designs* 21 (2013), 232-252.
- [9] C. Godsil, Compact graphs and equitable partitions, *Linear Algebra and Its Application* 255(1997), 259-266.

Email: alaeiyan@iust.ac.ir;

Email: zahra.shokoohi75@yahoo.com